



# Welcome to Verzenio

  
**Verzenio**<sup>®</sup>  
abemaciclib  
50 | 100 | 150 | 200 mg tablets

# Table of contents

## **ABOUT VERZENIO..... 3**

What is Verzenio?  
How Verzenio is given  
How Verzenio works  
How is Verzenio different?

## **DOSING..... 5**

## **VERZENIO + AN AROMATASE INHIBITOR (AI)..... 7**

Effectiveness  
Side effects

## **VERZENIO + FULVESTRANT..... 9**

Effectiveness  
Side effects

## **VERZENIO BY ITSELF..... 11**

Effectiveness  
Side effects

## **ADDRESSING SIDE EFFECTS..... 13**

Addressing diarrhea  
Immediate 24-hour action plan

## **SAVINGS & SUPPORT..... 15**

Verzenio Continuous Care™ Program  
Savings Card

## **INDICATION AND SAFETY SUMMARY..... 17**

Personalized Care  
Real Patient Stories



**Personalized support that's here for you**

*Find out how to enroll on [page 15](#)*

The Verzenio Continuous Care Team is not made up of medical professionals.  
Your doctor is your source for medical advice.

# About Verzenio

Whether you're newly diagnosed or just starting treatment, this brochure can help you learn more about Verzenio.

## Make the most of every day with everyday Verzenio

\*150 mg or 200 mg orally twice a day, as directed by your doctor.

**Verzenio is a prescription medicine used to treat a type of breast cancer in adults. It is a medicine:**

- You can take it if you have a type of breast cancer called HR+, HER2– (hormone receptor positive, human epidermal growth factor receptor 2 negative) and the cancer has spread to other parts of the body (metastasized)
- That is given with an aromatase inhibitor as initial endocrine-based therapy. If the cancer has gotten worse after hormone therapy, then Verzenio is given with fulvestrant. If the cancer has gotten worse after both hormone therapy and chemotherapy, then Verzenio is given by itself

It is not known if Verzenio is safe and effective in children.

### SELECT SAFETY INFORMATION

#### Warnings - Verzenio may cause serious side effects, including:

Diarrhea is common with Verzenio, may sometimes be severe and may cause dehydration or infection. The most common time to develop diarrhea is during the first month of Verzenio treatment. If you develop diarrhea during treatment with Verzenio, your healthcare provider may tell you to temporarily stop taking it, stop your treatment, or decrease your dose.

If you have any loose stools, start taking an antidiarrheal medicine (such as loperamide), drink more fluids, and tell your healthcare provider right away.

Please see Indication and Safety Summary on pages [17-18](#).

## What is Verzenio?

Verzenio is a **targeted treatment** known as a **CDK4 & 6 inhibitor**. Verzenio is **not chemotherapy**. **Verzenio is a prescription medicine used to treat a type of breast cancer in adults**. It is a medicine you can take if you have a type of breast cancer called **HR+, HER2-** and the cancer has **spread to other parts of the body (metastasized)**.

## How Verzenio is given

- with an aromatase inhibitor as initial endocrine-based therapy
- if the cancer has gotten worse after hormone therapy, then Verzenio is given with fulvestrant
- if the cancer has gotten worse after both hormone therapy and chemotherapy, then Verzenio is given by itself

It is not known if Verzenio is safe and effective in children.

CDK4 & 6=cyclin-dependent kinases 4 and 6.

HER2-=human epidermal growth factor receptor 2 negative.

HR+=hormone receptor positive.

## How Verzenio works

**Verzenio targets breast cancer cells where they grow and divide**

- Verzenio is in a class of drugs known as CDK4 & 6 inhibitors. CDK4 & 6 are proteins that control how fast cells grow and divide
- These proteins are found on both normal and cancer cells. They become overactive in metastatic breast cancer, causing cells to grow and divide uncontrollably. This leads to the spread of cancer
- Verzenio interrupts these proteins and cells just as they are deciding to grow and divide. It slows down cancer cell growth and division, causing cancer cells to become inactive or even die
- In preclinical studies, continuous exposure to Verzenio put the brakes on CDK4 & 6 proteins, which delayed cancer cells from growing and dividing

## How is Verzenio different?

**Verzenio is taken differently—it's the first and only CDK4 & 6 inhibitor that can be taken every day, twice a day\***

Verzenio is a CDK4 & 6 inhibitor. It is the first and only oral tablet of its kind that can be taken every day for the treatment of HR+, HER2- metastatic breast cancer.

\*150 mg or 200 mg orally twice a day, as directed by your doctor.

## SELECT SAFETY INFORMATION

**Warnings - Verzenio may cause serious side effects, including:**

Low white blood cell counts (neutropenia) are common with Verzenio and may cause serious infections that can lead to death. Your healthcare provider should check your white blood cell counts before and during treatment. If you develop low white blood cell counts during treatment with Verzenio, your healthcare provider may tell you to temporarily stop taking it, decrease your dose, or wait before starting your next month of treatment. Tell your healthcare provider right away if you have signs and symptoms of low white blood cell counts or infections, such as fever and chills.

**Please see Indication and Safety Summary on pages 17-18.**





# Dosing


Learn more about how to take Verzenio.

## VERZENIO + AN


### AROMATASE INHIBITOR (AI)\*


1 150-mg tablet by mouth in the morning 


1 150-mg tablet by mouth in the evening 

You will receive 4 individual packs,<sup>‡</sup> each containing your dose of 150-mg tablets for 7 days 


### VERZENIO + FULVESTRANT<sup>†</sup>


1 150-mg tablet by mouth in the morning 


1 150-mg tablet by mouth in the evening 

You will receive 4 individual packs,<sup>‡</sup> each containing your dose of 150-mg tablets for 7 days 

### VERZENIO BY ITSELF

1 200-mg tablet by mouth in the morning 

1 200-mg tablet by mouth in the evening 

You will receive 4 individual packs,<sup>‡</sup> each containing your dose of 200-mg tablets for 7 days 

**Try to establish a routine of taking your morning and evening doses at about the same time each day.**

- Do not stop taking Verzenio or change the dose without talking to your doctor
- Multiple tablet strengths (50 mg, 100 mg, 150 mg, and 200 mg) allow your doctor to make dose adjustments based on your tolerance
- If you need to reduce your dose due to tolerability, Verzenio is approved to be dosed as low as 50 mg twice daily, as directed by your doctor
- No matter what dose you are prescribed, you will take 1 tablet, 2 times a day, unless your doctor tells you otherwise

\*The recommended dose of an AI is 1 tablet taken by mouth once daily. The recommended doses for letrozole and anastrozole are 2.5 mg and 1 mg, respectively.

<sup>†</sup>The recommended starting dose of fulvestrant is 500 mg administered on days 1, 15, and 29 for the first month and once monthly thereafter.

<sup>‡</sup>Each of these packs includes 14 individual tablets.

## SELECT SAFETY INFORMATION

### Warnings - Verzenio may cause serious side effects, including:

Verzenio may cause severe or life-threatening inflammation (swelling) of the lungs during treatment that can lead to death. If you develop lung problems during treatment with Verzenio, your healthcare provider may tell you to temporarily stop taking it, decrease your dose, or stop your treatment. Tell your healthcare provider right away if you have any new or worsening symptoms, including:

- Trouble breathing or shortness of breath
- Cough with or without mucus
- Chest pain

Please see Indication and Safety Summary on pages [17-18](#).

## What should I do if I miss a dose or get sick after taking a dose?

- If you miss a dose of Verzenio or get sick after taking a dose, take your next dose at your regular time. Do not take an extra dose of Verzenio to make up for any dose you may have missed
- For example, if you forget to take your morning dose, then you should only take your regular evening dose rather than doubling it



---

If you take more Verzenio than was prescribed, call your doctor or go to the nearest hospital emergency room right away

---

## How should I take Verzenio?

- Swallow the prescribed number of Verzenio tablets whole. Do not chew, crush, or split before swallowing
- You can take Verzenio with or without food
- Avoid grapefruit and grapefruit products while taking Verzenio. Grapefruit may increase the amount of Verzenio in your blood

## Can I take other medicines with Verzenio?

- Verzenio and certain other medicines can affect each other, causing serious side effects or changing the effect of Verzenio or other medications
- Before starting any medication, make sure you talk to your doctor about all your other medications. Some examples include prescription and over-the-counter medicines, vitamins, and herbal supplements. Your doctor may need to change your dose

If you have questions about Verzenio or need more information on how to take it, you should talk to your doctor. You can also call us at 1-844-VERZENIO (1-844-837-9364).

### SELECT SAFETY INFORMATION

#### Warnings - Verzenio may cause serious side effects, including:

Verzenio can cause serious liver problems. Your healthcare provider should do blood tests to check your liver before and during treatment. If you develop liver problems during treatment with Verzenio, your healthcare provider may reduce your dose or stop your treatment. Tell your healthcare provider right away if you have any of the following signs or symptoms of liver problems:

- Feeling very tired
- Pain on the upper right side of your stomach area (abdomen)
- Loss of appetite
- Bleeding or bruising more easily than normal

Please see Indication and Safety Summary on pages [17-18](#).

# Verzenio + an aromatase inhibitor (AI)

Explore the effectiveness and side effects of Verzenio + an AI.

Verzenio was studied in combination with an AI as initial endocrine-based therapy in 493 postmenopausal women with newly diagnosed, hormone receptor positive (HR+), human epidermal growth factor receptor 2 negative (HER2-) advanced or metastatic breast cancer. In the study, 328 patients were treated with Verzenio + an AI and 165 were treated with an AI alone.



## LIVE LONGER WITHOUT CANCER GETTING WORSE

Verzenio + an AI delayed disease progression for over 2 years (a median of 28.2 months) vs 14.8 months with an AI alone



## More than half of the women saw their **tumors shrink**

In a clinical study, 55.4% of women on Verzenio + an AI saw their tumors shrink vs 40.2% on an AI alone.

### COMPLETE RESPONSE:

#### NO DETECTABLE SIGNS OF CANCER

3.4% of women on Verzenio + an AI had a complete response vs 0% with an AI alone.

### PARTIAL RESPONSE:

#### TUMOR SHRINKAGE OF 30% OR MORE

52.1% of women on Verzenio + an AI had a partial response vs 40.2% with an AI alone.

Clinical trials are still ongoing to determine if there is an overall survival benefit.

## SELECT SAFETY INFORMATION

### Verzenio may cause serious side effects, including:

Verzenio may cause blood clots in your veins, or in the arteries of your lungs. Verzenio may cause serious blood clots that have led to death. If you develop blood clots during treatment with Verzenio, your healthcare provider may tell you to temporarily stop taking it. Tell your healthcare provider right away if you have any of the following signs and symptoms of a blood clot:

- Pain or swelling in your arms or legs
- Shortness of breath
- Chest pain
- Fast breathing
- Fast heart rate


Please see Indication and Safety Summary on pages [17-18](#).

## Side effects

### Most common side effects

- Diarrhea
- Low white blood cell count (leukopenia/neutropenia)
- Infections
- Nausea
- Low red blood cell count (anemia)
- Vomiting
- Hair loss or thinning (alopecia)
- Decreased appetite
- Tiredness
- Abdominal pain

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of Verzenio. For more information, ask your doctor or pharmacist.



---

TO LEARN MORE ABOUT ADDRESSING  
SIDE EFFECTS, TURN TO PAGES [13-14](#)

---

### SELECT SAFETY INFORMATION

#### Verzenio may cause serious side effects, including:

Verzenio can harm your unborn baby. Use effective birth control during treatment and for 3 weeks after the last dose of Verzenio and do not breastfeed during treatment with Verzenio and for at least 3 weeks after your last dose. Verzenio may affect the ability of males to father a child.

Please see Indication and Safety Summary on pages [17-18](#).



# Verzenio + fulvestrant

Explore the effectiveness and side effects of Verzenio + fulvestrant.

Verzenio was studied in a clinical trial of 669 women with hormone receptor positive (HR+), human epidermal growth factor receptor 2 negative (HER2-) metastatic breast cancer, whose disease had progressed after hormone treatment: 446 patients were treated with Verzenio + fulvestrant, and 223 were treated with fulvestrant alone.




## WOMEN LIVED LONGER ON VERZENIO + FULVESTRANT

In the clinical study, women taking Verzenio + fulvestrant lived for a median of 46.7 months vs 37.3 months on fulvestrant alone



## MORE TIME WITHOUT DISEASE PROGRESSION

In the same study, Verzenio + fulvestrant delayed disease progression for a median of 16.4 months vs 9.3 months with fulvestrant alone



## TWICE AS MANY WOMEN SAW THEIR TUMORS SHRINK

48.1% of patients on Verzenio + fulvestrant saw their tumors shrink vs 21.3% with fulvestrant alone

### COMPLETE RESPONSE: NO DETECTABLE SIGNS OF CANCER

In a clinical study, 3.5% of women on Verzenio + fulvestrant had a complete response vs 0% with fulvestrant alone.

### PARTIAL RESPONSE: TUMOR SHRINKAGE OF 30% OR MORE

In the same study, 44.7% of women on Verzenio + fulvestrant had a partial response vs 21.3% with fulvestrant alone.

## SELECT SAFETY INFORMATION

### Common side effects

The most common side effects of Verzenio include:

- Nausea
- Infections
- Low red blood cell counts (anemia)
- Decreased appetite
- Headache
- Hair thinning or hair loss (alopecia)
- Abdominal pain
- Tiredness
- Low white blood cell counts (leukopenia)
- Vomiting
- Low platelet counts (thrombocytopenia)

These are not all the possible side effects of Verzenio

Please see Indication and Safety Summary on pages [17-18](#).

## Side effects

### Most common side effects


- Diarrhea
- Nausea
- Infections
- Low white blood cell count (leukopenia/neutropenia)
- Abdominal pain
- Low red blood cell count (anemia)
- Tiredness
- Decreased appetite
- Vomiting
- Headache

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of Verzenio. For more information, ask your doctor or pharmacist.

---

TO LEARN MORE ABOUT ADDRESSING  
SIDE EFFECTS, TURN TO PAGES [13-14](#)

---



### SELECT SAFETY INFORMATION

#### Before using

Before you use Verzenio, tell your healthcare provider about all your medical conditions, including:

- If you have fever, chills, or other signs of infection.
- If you have a history of blood clots in your veins.
- Have lung or breathing problems.
- Have liver or kidney problems.
- If you are pregnant, plan to become pregnant, or are breastfeeding.
- About all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take a medicine that contains ketoconazole.

Please see Indication and Safety Summary on pages [17-18](#).

# Verzenio by itself

Explore the effectiveness and side effects of Verzenio by itself.

Verzenio was studied in 132 adult patients with hormone receptor positive (HR+), human epidermal growth factor receptor 2 negative (HER2-) advanced or metastatic breast cancer who had progressed on or after prior hormone treatments, a taxane, and 1 or 2 chemotherapy regimens in the metastatic setting. All patients were treated with only Verzenio.

## VERZENIO SHRINKS TUMORS AND DELAYS THE SPREAD OF CANCER

When other advanced or metastatic breast cancer treatments (hormonal treatments and prior chemotherapy) stopped working, Verzenio helped delay the growth and spread of cancer in a clinical study.

## TUMOR

SHRINKAGE OF  
**30% or more**

19.7% of women taking Verzenio alone saw their tumors shrink by 30% or more. In the same study, an independent review found similar results (17.4% of women)

---

LEARN MORE ABOUT VERZENIO BY ITSELF AT [VERZENIO.COM](https://www.verzenio.com)

---

## SELECT SAFETY INFORMATION

### Warnings - Verzenio may cause serious side effects, including:

Diarrhea is common with Verzenio, may sometimes be severe and may cause dehydration or infection. The most common time to develop diarrhea is during the first month of Verzenio treatment. If you develop diarrhea during treatment with Verzenio, your healthcare provider may tell you to temporarily stop taking it, stop your treatment, or decrease your dose.

If you have any loose stools, start taking an antidiarrheal medicine (such as loperamide), drink more fluids, and tell your healthcare provider right away.


Please see Indication and Safety Summary on pages [17-18](#).

## Side effects

### Most common side effects

- Diarrhea
- Tiredness
- Nausea
- Decreased appetite
- Abdominal pain
- Low white blood cell count (neutropenia)
- Vomiting
- Infections
- Low red blood cell count (anemia)
- Headache
- Low platelet count (thrombocytopenia)

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of Verzenio. For more information, ask your doctor or pharmacist.



---

TO LEARN MORE ABOUT ADDRESSING  
SIDE EFFECTS, TURN TO PAGES [13-14](#)

---

### SELECT SAFETY INFORMATION

#### Warnings - Verzenio may cause serious side effects, including:

Low white blood cell counts (neutropenia) are common with Verzenio and may cause serious infections that can lead to death. Your healthcare provider should check your white blood cell counts before and during treatment. If you develop low white blood cell counts during treatment with Verzenio, your healthcare provider may tell you to temporarily stop taking it, decrease your dose, or wait before starting your next month of treatment. Tell your healthcare provider right away if you have signs and symptoms of low white blood cell counts or infections, such as fever and chills.

Please see Indication and Safety Summary on pages [17-18](#).

# Addressing side effects

Learn how to address some common side effects.

Verzenio may cause side effects, including diarrhea, low white blood cell counts, severe inflammation of the lungs, liver problems, or blood clots. Verzenio can harm your unborn baby.

These are not all the possible side effects of Verzenio. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call **1-800-FDA-1088**.

You may also report side effects to Eli Lilly and Company at **1-800-545-5979**.

## Addressing diarrhea

The first step to managing side effects is to know what to expect.

Diarrhea is one of the most common side effects with Verzenio. Read on to learn more about how to **EXPECT, PLAN, and ACT** if you experience diarrhea while taking Verzenio.

Diarrhea can be managed with the appropriate steps, so **EXPECT, PLAN, and ACT**.

### EXPECT: DIARRHEA IS A COMMON SIDE EFFECT

Diarrhea is a common side effect with Verzenio and may sometimes be severe, but you and your doctor can take steps to manage it. If diarrhea occurs, it will most likely happen in the first week. You should be on the lookout for cramping or frequent, loose, or watery stools.

### PLAN: PLAN AHEAD OF TIME

It is important to plan ahead of time with your doctor. Make sure you have an over-the-counter anti-diarrheal medicine, such as loperamide (also known as Imodium®), available at home before you start taking Verzenio.

### ACT: IMMEDIATE 24-HOUR ACTION PLAN

If you begin experiencing diarrhea, it is very important that you take action immediately at the first signs of symptoms. The best way to manage diarrhea is to start an antidiarrheal at the first sign and call your doctor, who may make a dose change.

## SELECT SAFETY INFORMATION

### Warnings - Verzenio may cause serious side effects, including:

Verzenio may cause severe or life-threatening inflammation (swelling) of the lungs during treatment that can lead to death. If you develop lung problems during treatment with Verzenio, your healthcare provider may tell you to temporarily stop taking it, decrease your dose, or stop your treatment. Tell your healthcare provider right away if you have any new or worsening symptoms, including:

- Trouble breathing or shortness of breath
- Cough with or without mucus
- Chest pain

Please see Indication and Safety Summary on pages [17-18](#).

# Immediate 24-hour action plan



### STEP 1: Start an antidiarrheal and call your doctor

If you are experiencing diarrhea, you need to start taking an antidiarrheal as directed on the packaging and call your doctor or nurse immediately so they are aware and can help you manage the diarrhea.



### STEP 2: Stay hydrated and drink clear fluids

Drink at least 8-12 glasses of clear fluids in the first 24 hours you experience symptoms (and continue any day you experience diarrhea) to avoid dehydration.



### STEP 3: Watch for improvement and follow up with your doctor

Ideally, the antidiarrheal will be able to help resolve your symptoms, but **if symptoms continue after the first 24 hours, you need to talk to your doctor or nurse that day.** He or she may consider adjusting your dose, with the ultimate goal of finding which dose is right for you. Verzenio is effective at different doses, so it is important to find what dose is right for your body.



### ADDITIONAL MONITORING MONITORING BLOOD COUNTS AND LIVER FUNCTION

Some patients experienced low white blood cell counts (neutropenia) and/or liver problems. Your doctor will do blood tests to monitor your blood cell counts and to check for liver problems before and during treatment.

## SELECT SAFETY INFORMATION

### Warnings - Verzenio may cause serious side effects, including:

Verzenio can cause serious liver problems. Your healthcare provider should do blood tests to check your liver before and during treatment. If you develop liver problems during treatment with Verzenio, your healthcare provider may reduce your dose or stop your treatment. Tell your healthcare provider right away if you have any of the following signs or symptoms of liver problems:

- Feeling very tired
- Pain on the upper right side of your stomach area (abdomen)
- Loss of appetite
- Bleeding or bruising more easily than normal

Please see Indication and Safety Summary on pages [17-18](#).



# Savings & support

See how you can save on Verzenio and get support along the way.

## Personalized care that's here for you

### Verzenio Continuous Care™



We created the Verzenio Continuous Care Program to give you personalized support and help you have a positive experience while taking Verzenio. Your Verzenio Continuous Care Team will serve as your dedicated partner throughout every stage of your treatment.

The Verzenio Continuous Care Team is not made up of medical professionals. Your doctor is your source for medical advice.

Through 1-on-1 check-ins, your Verzenio Continuous Care Team can support you by:

- Helping you get started on Verzenio quickly
- Focusing on the insurance process, so you can focus on treatment
- Being there to answer specific questions or just lend an ear

## 3 simple ways to enroll

**1**

Enroll at  
**verzenio.com**

OR

**2**

Call  
**1-844-VERZENIO**  
(1-844-837-9364)

OR

**3**

Ask a member of  
your healthcare team  
about signing up

## Savings Card

Paying for treatment shouldn't be an additional concern for you and your loved ones, so we've created the Verzenio Savings Card,\* which may help you manage treatment costs.



---

**Scan to learn more about  
how to save on Verzenio**

---

\*Governmental beneficiaries excluded, terms and conditions apply.

---

Visit [Verzenio.com/savings-support](https://www.verzenio.com/savings-support) for more information  
on the Verzenio Savings Card

---



## INDICATION AND SAFETY SUMMARY

Verzenio® (ver-ZEN-ee-oh) is used to treat certain types of breast cancer known as HR+/HER2– (hormone receptor positive/human epidermal growth factor receptor 2 negative) breast cancer.

It is a medicine you can take if:

- The cancer has spread to other parts of the body (metastasized). If you are female or male, then Verzenio is given with an aromatase inhibitor as initial endocrine-based therapy. If you are a female or male, and the cancer has gotten worse after hormonal therapy, then Verzenio is given with fulvestrant. If you are female or male and the cancer has gotten worse after both hormonal therapy and chemotherapy, then Verzenio is given by itself.

It is not known if Verzenio is safe and effective in children.

### Warnings - Verzenio may cause serious side effects, including:

Diarrhea is common with Verzenio, may sometimes be severe and may cause dehydration or infection. The most common time to develop diarrhea is during the first month of Verzenio treatment. If you develop diarrhea during treatment with Verzenio, your healthcare provider may tell you to temporarily stop taking it, stop your treatment, or decrease your dose.

If you have any loose stools, start taking an antidiarrheal medicine (such as loperamide), drink more fluids, and tell your healthcare provider right away.

Low white blood cell counts (neutropenia) are common with Verzenio and may cause serious infections that can lead to death. Your healthcare provider should check your white blood cell counts before and during treatment. If you develop low white blood cell counts during treatment with Verzenio, your healthcare provider may tell you to temporarily stop taking it, decrease your dose, or wait before starting your next month of treatment. Tell your healthcare provider right away if you have signs and symptoms of low white blood cell counts or infections, such as fever and chills.

Verzenio may cause severe or life-threatening inflammation (swelling) of the lungs during treatment that can lead to death. If you develop lung problems during treatment with Verzenio, your healthcare provider may tell you to temporarily stop taking it, decrease your dose, or stop your treatment. Tell your healthcare provider right away if you have any new or worsening symptoms, including:

- Trouble breathing or shortness of breath
- Cough with or without mucus
- Chest pain

Verzenio can cause serious liver problems. Your healthcare provider should do blood tests to check your liver before and during treatment. If you develop liver problems during treatment with Verzenio, your health care provider may reduce your dose or stop your treatment. Tell your healthcare provider right away if you have any of the following signs or symptoms of liver problems:

- Feeling very tired
- Pain on the upper right side of your stomach area (abdomen)
- Loss of appetite
- Bleeding or bruising more easily than normal

Verzenio may cause blood clots in your veins, or in the arteries of your lungs. Verzenio may cause serious blood clots that have led to death. If you develop blood clots during treatment with Verzenio, your healthcare provider may tell you to temporarily stop taking it. Tell your healthcare provider right away if you have any of the following signs and symptoms of a blood clot:

- Pain or swelling in your arms or legs
- Shortness of breath
- Chest pain
- Fast breathing
- Fast heart rate

## INDICATION AND SAFETY SUMMARY (continued)

Verzenio can harm your unborn baby. Use effective birth control (contraception) during treatment and for 3 weeks after the last dose of Verzenio and do not breastfeed during treatment with Verzenio and for at least 3 weeks after your last dose. Verzenio may affect the ability of males to father a child.

### Common side effects

The most common side effects of Verzenio include:

- Nausea
- Decreased appetite
- Abdominal pain
- Vomiting
- Infections
- Headache
- Tiredness
- Low platelet counts (thrombocytopenia)
- Low red blood cell counts (anemia)
- Hair thinning or hair loss (alopecia)
- Low white blood cell counts (leukopenia)

These are not all the possible side effects of Verzenio.

Tell your healthcare provider if you have any side effects. **You can report side effects at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### Before using

Before you use Verzenio, tell your healthcare provider about all your medical conditions, including:

- If you have fever, chills, or other signs of infection.
- If you are pregnant, plan to become pregnant, or are breastfeeding.
- If you have a history of blood clots in your veins.
- About all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take a medicine that contains ketoconazole.
- Have lung or breathing problems.
- Have liver or kidney problems.

### How to take

- Take Verzenio exactly as your healthcare provider tells you.
- Swallow Verzenio tablets whole. Do not chew, crush, or split the tablets before swallowing. Do not take Verzenio tablets if they are broken, cracked, or damaged.
- Your healthcare provider may change your dose if needed. Do not stop taking Verzenio or change the dose without talking to your healthcare provider.
- Take your doses of Verzenio at about the same time every day.
- Verzenio may be taken with or without food.
- If you vomit or miss a dose of Verzenio, take your next dose at your regular time. Do not take 2 doses at the same time to make up for the missed dose.

### What to avoid during treatment

- Avoid taking ketoconazole during treatment with Verzenio. Tell your healthcare provider if you take a medicine that contains ketoconazole.
- Avoid grapefruit and products that contain grapefruit during treatment with Verzenio. Grapefruit may increase the amount of Verzenio in your blood.

### Learn more

Verzenio is a prescription medicine. For more information, call 1-800-545-5979 or go to [verzenio.com](http://verzenio.com).

This summary provides basic information about Verzenio but does not include all information known about this medicine. Read the information that comes with your prescription each time your prescription is filled. This information does not take the place of talking with your healthcare provider. Be sure to talk to your healthcare provider about Verzenio and how to take it. Your healthcare provider is the best person to help you decide if Verzenio is right for you.

AL CON BS\_MBC 03MAR2023

Verzenio® is a registered trademark owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.



To everyone living with  
HR+, HER2- metastatic breast cancer  
and taking Verzenio, we're with you

**Personalized care that's here for you**

Whether you have questions or just need someone to talk to, Verzenio  
Continuous Care is here for you.

The Verzenio Continuous Care Team is not made up of medical professionals.  
Your doctor is your source for medical advice.

---

**Sign up for Verzenio Continuous Care today**

---


[verzenio.com/savings-support/continuous-care](https://verzenio.com/savings-support/continuous-care)  
**1-844-VERZENIO (1-844-837-9364)**

  
**Verzenio**<sup>®</sup>  
abemaciclib  
50 | 100 | 150 | 200 mg tablets

Hear inspiring  
stories from real,  
relentless women

Visit [verzenio.com](https://www.verzenio.com)

Learn about Verzenio,  
advocacy groups, and  
so much more.



---

Visit the Verzenio Spanish website  
at [verzenio.com/es](https://www.verzenio.com/es)

---

Verzenio<sup>®</sup> is a registered trademark and Verzenio Continuous Care<sup>™</sup> is a trademark owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates. Other product/company names mentioned herein are the trademarks of their respective owners.

PP-AL-US-4134 11/2023 ©Lilly USA, LLC 2023.  
All rights reserved.



*Lilly*